



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

SOLUTION BY THE PROPOSER.

Let a = number of letters and spaces in heading.

b = number of spaces on 80 scale of right stop.

Then $\frac{80 - a}{2}$ = number on 80 scale to begin heading. But by the method used this number is found as $\left\{ \left(40 - \frac{b}{2} \right) - \frac{a}{2} \right\} + \frac{b}{2}$.


GEOMETRY.

418. Proposed by MRS. H. E. TREFETHEN, Waterville, Maine.

The difference of the squares of the two interior diagonals of a cyclic quadrilateral is to twice their rectangle as the distance between their mid-points is to the third diagonal.

SOLUTION BY C. N. SCHMALL, New York, N. Y.

Let $ABCD$ be a cyclic quadrilateral. Let the sides BA, CD meet in P , and the sides DA, CB in Q . Draw PQ . The resulting figure is a *complete quadri-*



lateral and PQ is the *third diagonal*. Let L, M, N , be the mid-points of the three diagonals. Then we have to prove that

$$\frac{AC^2 - BD^2}{2AC \cdot BD} = \frac{MN}{PQ}.$$

It is well known that the points, L, M, N , are collinear. (See RUSSELL'S *Sequel to Elem. Geom.*, p. 154, ex. 5; or LACHLAN'S *Pure Geom.*, p. 93, § 151; or MILNE'S *Cross-Ratio Geom.*, p. 113, ex. 3.) Hence $MN = LN - LM$, or

$$\frac{MN}{PQ} = \frac{LN}{PQ} - \frac{LM}{PQ}. \quad (1)$$

Draw QN and prolong it to R , making $NR = QN$. Draw AR . Then $QCRA$

is a parallelogram. Hence QC is parallel to AR and

$$QC = AR. \quad (2)$$

Now, the triangles PAC, PDB , are similar; for $\angle PCA = \angle PBD$, both standing on the arc AD in a cyclic quadrilateral. Hence

$$AC : DB :: PA : PD. \quad (3)$$

Likewise, the triangles QCA, QDB are similar; and

$$AC : DB :: QC : QD. \quad (4)$$

From (3) and (4),

$$PA : PD :: QC : QD;$$

or

$$PD : QD :: PA : QC;$$

or, by (2),

$$PD : QD :: PA : AR. \quad (5)$$

Again,

$$\angle PDQ = \angle PAR, \quad (6)$$

for

$$\angle PDQ = 180^\circ - \angle ADC = \angle ABC$$

and

$$\angle PAR = \angle ABC.$$

Hence, by (5) and (6) the triangles PDQ and PAR are similar; since they have an angle in each equal and the including sides proportional. Therefore,

$$PA : PD :: PR : PQ;$$

or

$$PA : PD :: 2LN : PQ. \quad (7)$$

By (3) and (7) we have

$$AC : BD :: 2LN : PQ. \quad (8)$$

In a similar way, by drawing QM and prolonging it to S (say), making $MS = QM$, and then drawing PS , we can show that

$$AC : BD :: PQ : 2LM. \quad (9)$$

From (8) and (9),

$$\frac{LN}{PQ} = \frac{1}{2} \frac{AC}{BD}, \quad \frac{LM}{PQ} = \frac{1}{2} \frac{BD}{AC}.$$

Substituting these values in (1), we have

$$\frac{MN}{PQ} = \frac{1}{2} \left(\frac{AC}{BD} - \frac{BD}{AC} \right); \quad \text{or} \quad \frac{MN}{PQ} = \frac{\overline{AC}^2 - \overline{BD}^2}{2AC \cdot BD}.$$

CALCULUS.

333. Proposed by C. N. SCHMALL, New York City.

Evaluate $\int \int \int \sqrt{\frac{1 - (x^2 + y^2 + z^2)}{1 + x^2 + y^2 + z^2}} dx dy dz$, where $x^2 + y^2 + z^2 < 1$.